

## A34

### **Protein kinase C- $\zeta$ is involved in the inhibition of eosinophil migration**

Eva M Sturm, Gerald P Parzmair, Petra Luschnig-Schratl, Viktória Kónya and Ákos Heinemann

*Institute of Experimental and Clinical Pharmacology, Medical University of Graz, 8010 Graz, Austria*

*E-mail: akos.heinemann@medunigraz.at*

#### **Background**

The accumulation of eosinophils in lung tissue is a hallmark of asthma, and it is believed that eosinophils play a crucial pathogenic role in allergic inflammation. Therefore, eosinophils are currently considered a major therapeutic target in allergic diseases. Prostaglandin (PG) E<sub>2</sub> exerts anti-inflammatory and broncho-protective mechanisms in asthma, but the underlying mechanisms have remained unclear. We have shown previously that PGE<sub>2</sub> and the EP<sub>2</sub> receptor agonist butaprost inhibit eosinophil trafficking *in vitro* and *in vivo* [1].

#### **Methods**

Human eosinophils were purified by negative magnetic selection from peripheral blood. Chemotaxis was determined in 48-well microBoyden chambers and migrated eosinophils were enumerated by flow cytometry.

#### **Results**

The chemotaxis of human eosinophils towards the chemoattractant eotaxin was attenuated by PGE<sub>2</sub> and the selective EP<sub>4</sub> agonist ONO-AE1-329 in a concentration-dependent manner. Pretreatment of eosinophils with the adenylyl cyclase inhibitor SQ-22536, the protein kinase A inhibitor H-89 or the p38 MAP kinase inhibitor SB-202190 did not prevent the inhibitory effect of PGE<sub>2</sub>, while the phosphatidyl inositol 3-kinase (PI3K) inhibitor LY-294002, triciribine, a specific inhibitor of Akt phosphorylation, and a myristoylated pseudosubstrate of protein kinase (PK) C- $\zeta$  (mPS), partially or completely reversed the inhibitory effect of PGE<sub>2</sub> and ONO-AE1-329 on the migration of eosinophils towards eotaxin.

#### **Conclusions**

Protein kinase C is an increasingly diverse family of enzymes that has been implicated in a range of cellular functions within the eosinophil. The present data show that the PI3K/Akt/PKC- $\zeta$  pathway is involved in the negative regulation of the migration of human eosinophils mediated by EP<sub>2</sub> and EP<sub>4</sub> receptor activation.

#### **Reference**

1. Sturm EM, Schratl P, Schuligoi R, Konya V, Sturm GJ, Lippe IT, Peskar BA, Heinemann A: **Prostaglandin E<sub>2</sub> inhibits eosinophil trafficking through E-prostanoid 2 receptors.** *J Immunol* 2008, **181**:7273–7283.