

A64

Endothelin-1 increases osteoclastic bone resorption via endothelin A receptors during orthodontic tooth movement in rats

Špela Sprogar^{1,2}, Alja Meh³, Tomaž Vaupotič⁴, Andrej Cör⁵, Martina Drevenšek³ and Gorazd Drevenšek¹

¹*Institute of Pharmacology and Experimental Toxicology, Faculty of Medicine, University of Ljubljana, 1000 Ljubljana, Slovenia*

²*Ort-line, Institute for Orthodontics and General Dentistry, 1000 Ljubljana, Slovenia*

³*Department of Orthodontics, Faculty of Medicine, University of Ljubljana, 1000 Ljubljana, Slovenia*

⁴*Institute of Biochemistry, Faculty of Medicine, University of Ljubljana, 1000 Ljubljana, Slovenia*

⁵*Department of Medical and Natural Sciences, College of Health Care, University of Primorska, 6000 Koper, Slovenia*

E-mail: ssprogar@gmail.com

Background

The involvement of the endothelin signaling system during orthodontic tooth movement has not been explained yet. Therefore, the aim of this study was to determine the role of endothelins ET-1, ET-2 and ET-3 and both receptor subtypes ET_A and ET_B during all the three phases of orthodontic tooth movement in a rat model.

Methods

The study was performed on male Wistar rats (n = 85). Orthodontic tooth movement was induced by a closed coil spring (F = 25 cN), which was placed between the upper left first molar and the upper incisors. The effects of the endothelin system were investigated using tezosentan, a non-selective endothelin antagonist, and TBC3214, a highly selective ET_A antagonist. Measurements of the distance between the upper left first molar and the ipsilateral incisor were performed on a weekly basis for 6 consecutive weeks. After that, the animals were sacrificed and tissue samples of the maxilla were taken for further biochemical and histological evaluations.

Results

Tezosentan increased tooth movement (p < 0.01). The opposite effect was shown using TBC3214, which decreased tooth movement (p < 0.01). On day 14, gene expression levels for ET-1 (p < 0.05) and ET-3 (p < 0.001) were increased compared to day 0. On day 28, a down-regulation of ET-3 was observed when compared to day 0 (p < 0.001). On day 42, ET-1 (p < 0.001) and ET-3 (p < 0.01) gene expression levels were strongly up-regulated, while ET-2 gene expression level was down-regulated (p < 0.01) when compared with day 0. The immunoreactivity of ET_A and ET_B significantly decreased on day 14 (p < 0.001) and increased on day 28 (p < 0.001). Alveolar bone volume was significantly higher in the TBC3214 group compared to the appliance only group (p < 0.001). Osteoclast volume was significantly lower in the TBC3214 group compared to the appliance only group (p < 0.05).

Conclusions

ET-1 and ET-3 are the endothelin isopeptides, which are involved in all three phases of orthodontic tooth movement. However, ET-1 is the predominant physiological form functioning during the late phase of orthodontic tooth movement. Gene and protein expression levels indicate that the major signaling pathway during the late phase of orthodontic tooth movement mainly involves ET_A receptors. During this phase ET-1 increases osteoclastic bone resorption via ET_A.